

## MEETING ABSTRACT

## Open Access

# The prevalence of ESBL-microorganisms among students of Southeast Asia living in Kazakhstan

Svetlana Kolesnichenko

From 4th International Conference for Healthcare and Medical Students (ICHAMS) 2014  
Dublin, Ireland. 24-25 October 2014

## Background

There are more and more information about ESBL-producing organisms wide spreading in Europe and Southeast Asia in recent years [1]. A large amount of students from Southeast Asia are staying in Kazakhstan, which raises the question about the distribution of microorganisms and aggravation of epidemiological situation. In our study we investigated the prevalence of ESBL-producing microorganisms among Southeast Asian students studying in Kazakhstan.

## Methods

Material for the study – feces. 139 fecal samples from foreign students from India and Pakistan were investigated in 2012 and 291 samples in 2013. The samples were examined for the presence of ESBL-producing in a medium containing ceftriaxone (6 mkg/ml). Identification of isolated microorganisms was performed by using MALDI-TOF spectrometer Microflex and software complex Biotyper of Bruker Daltoniks. Confirmation of ESBL-production was performed by double-discs method. Sensitivity to antibiotics was determined by disk diffusion method.

## Results

In 2012, as a result of screening plates with ceftriaxone from 139 stool samples 48 strains were cultivated. Those are 35 (72.9%) *Escherichia coli* strains and 1 (2%) strain of *Klebsiella pneumoniae*. After determining the sensitivity of these strains to antibiotics by disk diffusion method, ESBL-producing confirmed in 24 samples: 23 (47.9%) ESBL-positive *E. coli* strain and 1 (2%) strain of *K. pneumoniae*. According to the results of study in 2013, 280 microorganisms were cultivated from 292 stool samples, including 88.2% of *E. coli* (95% CI 20.24-34.14). Other

types of microorganisms were encountered in rare cases. 86% *E. coli* (240 strains) were ESBL-producers. Significant differences in the two methods: the method of double-discs and agar plates containing ceftriaxone was not obtained ( $p < 0.05$ ).

## Conclusions

The survey for 2012 was obtained 49.9% ESBL-producing strains, for 2013 - 86%, which means an increase in the prevalence of ESBL-producing strains. Using the agar plates containing ceftriaxone for ESBL screening significantly reduces the time and increases the level of studies of antibiotic-resistant strains of isolation.

Published: 27 October 2015

## References

1. Oteo J, Perez-Vazquez M, Campos J: Extended-spectrum [beta]-lactamase producing *Escherichia coli*: changing epidemiology and clinical impact. *Curr Opin Infect Dis* 2010, **23**(4):320-326.
2. Chaudhuri BN, Rodrigues C, Balaji V, Iyer R, Sekar U, Wattal C, et al: Incidence of ESBL producers amongst Gram-negative bacilli isolated from intra-abdominal infections across India. *J Assoc Physicians India* 2011, **59**:287-292.

doi:10.1186/1753-6561-9-S7-A28

**Cite this article as:** Kolesnichenko: The prevalence of ESBL-microorganisms among students of Southeast Asia living in Kazakhstan. *BMC Proceedings* 2015 **9**(Suppl 7):A28.

Faculty of General Medicine, Karganda State Medical University, Karganada, Kazakhstan



© 2015 Kolesnichenko This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.